

## Package leaflet: Information for the user

### Clindamycin 150 mg Capsules

clindamycin

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### **What is in this leaflet:**

1. What Clindamycin is and what it is used for
2. What you need to know before you take Clindamycin
3. How to take Clindamycin
4. Possible side effects
5. How to store Clindamycin
6. Contents of the pack and other information

### **1. What Clindamycin is and what it is used for**

Clindamycin 150 mg Capsules contain clindamycin hydrochloride which is an antibiotic used in the treatment of serious bacterial infections.

### **2. What you need to know before you take Clindamycin**

#### **Do not take Clindamycin**

If you are allergic (hypersensitive) to clindamycin, lincomycin or to any of the other ingredients in this medicine (listed in section 6).

#### **Warnings and precautions**

Talk to your doctor or pharmacist before using Clindamycin if:

- you have diarrhoea or usually get diarrhoea when you take antibiotics or have ever suffered from problems with your stomach or intestines. If you develop severe or prolonged or bloody diarrhoea during or after using this medicine, **tell your doctor immediately** since it may be necessary to interrupt the treatment. This may be a sign of bowel inflammation (pseudomembranous colitis) which can occur following treatment with antibiotics.
- you suffer from problems with your kidneys or liver.
- you suffer from asthma, eczema or hayfever.
- you develop any severe skin reactions or hypersensitivity to this medicine.

Acute kidney disorders may occur. Please inform your doctor about any medication you currently take and if you have any existing problems with your kidneys. If you experience decreased urine output, fluid retention causing swelling in your legs, ankles or feet, shortness of breath, or nausea you should contact your doctor immediately.

Due to the risk of inflammation of the oesophagus (oesophagitis) and oesophageal ulcer, it is important to always take Clindamycin Capsules exactly as described in section 3.

Clindamycin does not get into the brain and is therefore not suitable for treating serious infections in and around the brain. Your doctor may need to give you another antibiotic if you have these infections.

### **Other medicines and Clindamycin**

Tell your doctor or pharmacist or nurse if you are taking, have recently taken or might take any other medicines:

- muscle relaxants used for operations (neuromuscular blockers).
- oral contraceptive pills. You should use extra contraception such as condoms whilst taking this medicine and for seven days after taking it.
- warfarin or similar medicines – used to thin the blood. You may be more likely to have a bleed. Your doctor may need to take regular blood tests to check how well your blood can clot.
- CYP3A4 or CYP3A5 inducers like rifampicin may impact effectiveness of the medicine.

### **Pregnancy and breast-feeding**

#### **Pregnancy**

If you are pregnant or think you might be pregnant you should contact your doctor **before** taking Clindamycin. Ask your doctor or pharmacist for advice before taking any medicine.

#### **Breast-feeding**

Tell your doctor if you will be breast-feeding while taking this medicine as clindamycin may be passed into breast milk. Your doctor will decide if Clindamycin is appropriate for you. Although it is not likely that a nursing infant will take in very much of the active substance from the milk it drinks, if your baby gets bloodstained diarrhoea or shows any signs of illness, tell your doctor at once. You should stop breast-feeding if this happens.

### **Driving and using machines**

No effects on the ability to drive or use machines have been seen with Clindamycin.

### **Clindamycin contains lactose**

This medicine contains lactose, a type of sugar. If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicine.

## **3. How to take Clindamycin**

Always take this medicine exactly as your doctor has told you. Check with your doctor or pharmacist or nurse if you are not sure.

Clindamycin 150 mg Capsules should always be swallowed whole with a full glass of water, no less than 30 minutes before lying down, to avoid a possible irritation of the oesophagus.

### **Adults and Elderly Patients**

The recommended dose is between 150 and 450 mg (1 to 3 capsules) every 6 hours, depending on the severity of your infection.

### Use in children

This medicine is used for children who are able to swallow capsules. The recommended dose in children is between 12 and 25 mg/kg/day of bodyweight, divided into six hourly doses, depending on the severity of the infection. Clindamycin should be dosed based on total body weight regardless of obesity. Your doctor will work out the number of capsules that your child should have. If your child is unable to swallow capsules, talk to your doctor or pharmacist.

### Long-term use of Clindamycin

If you have to take this medicine for a long time, your doctor may arrange regular liver, kidney and blood tests. Do not miss these check-ups with your doctor. Long-term use can also make you more likely to get other infections that do not respond to Clindamycin treatment.

### If you take more Clindamycin than you should

If you accidentally take too many Clindamycin capsules, contact your doctor at once or go to the nearest hospital casualty department. Always take the labelled medicine package with you, whether there are any Clindamycin capsules left or not. Do not take any more capsules until your doctor tells you to.

### If you forget to take Clindamycin

If the forgotten dose is just a few hours late, use it straight away. If it is nearly time for your next dose miss out the forgotten one. **Do not take a double dose to make up for a missed dose.**

### If you stop taking Clindamycin

If you stop taking this medicine too soon your infection may come back again or get worse. Do not stop taking Clindamycin unless your doctor tells you to. If you have any further questions on how to take this product, ask your doctor or pharmacist.

## 4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Tell your doctor immediately** if you develop:

- severe, persistent or bloody diarrhoea (which may be associated with stomach pain or fever).  
This is an uncommon side effect which may occur during or after completing treatment with antibiotics and can be a sign of serious bowel inflammation or pseudomembranous colitis.
- signs of a severe allergic reaction such as sudden wheeziness, difficulty in breathing, dizziness, swelling of the eyelids or face or lips or throat or tongue, rash or itching (especially affecting the whole body).
- blistering and peeling of large areas of skin, fever, cough, feeling unwell and swelling of the gums, tongue or lips.
- yellowing of the skin and whites of the eyes (jaundice).
- potentially life-threatening skin rashes:
  - a widespread rash with blistering and peeling of large areas of skin, particularly around the mouth, nose, eyes or genitals, known as Stevens-Johnson syndrome, or a more severe form with extensive peeling of the skin (more than 30% of the body surface) known as *toxic epidermal necrolysis*
  - a rare skin eruption that is characterised by the rapid appearance of areas of red skin studded with small pustules (small blisters filled with white/yellow fluid) (Acute Generalised Exanthematous Pustulosis (AGEP))

- skin rash, which may blister, and looks like small targets (central dark spots surrounded by a paler area, with a dark ring around the edge - *erythema multiforme*)
  - widespread red skin rash with small pus-containing blisters (*bullous exfoliative dermatitis*)
  - fever, swollen lymph nodes or skin rash, these may be symptoms of a condition known as DRESS (Drug reaction with eosinophilia and systemic symptoms) and can be severe and life-threatening
- fluid retention causing swelling in your legs, ankles or feet, shortness of breath or nausea.

Other possible side effects may include:

**Common: may affect up to 1 in 10 people**

- abnormal liver function tests (poor liver function)
- pain in the stomach/abdomen, diarrhoea

**Uncommon: may affect up to 1 in 100 people**

- feeling sick or being sick
- rash characterized by a flat red area on the skin that is covered with small bumps, hives

**Not known: frequency cannot be estimated from the available data:**

- infection inside and around the vagina
- inflammation of the large intestine which causes abdominal pain, fever or diarrhoea due to infection by *Clostridioides difficile*
- effects on your blood system: reduced numbers of blood cells which may cause bruising or bleeding or weaken the immune system
- changes in the way things taste
- inflammation of the lining of the oesophagus (gullet)\*, open sores or lesions in the lining of the oesophagus (gullet)\*
- yellowing of the skin and whites of the eyes (jaundice)
- red or scaly skin (exfoliative dermatitis), red measles-like rash (rash morbilliform), itching

\* Side effects that are more likely to occur if Clindamycin Capsules are taken in a lying position and/or without enough water.

**Reporting of side effects**

If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard) or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Clindamycin**

- Keep this medicine out of the sight and reach of children.
- Do not use this medicine after the expiry date which is stated on the carton and blister foils after EXP. The expiry date refers to the last day of that month.
- Store below 25°C.

- Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

## **6. Contents of the pack and other information**

### **What Clindamycin contains**

The active substance is clindamycin hydrochloride. Each capsule contains clindamycin hydrochloride equivalent to 150 mg of clindamycin.

The other ingredients are lactose monohydrate, maize starch, talc, magnesium stearate.

Capsule: gelatin and titanium dioxide (E171).

Printing ink: shellac, soya lecithin, dimeticone (antifoam DC 1510), black iron oxide (E172).

### **What Clindamycin looks like and contents of the pack**

Clindamycin 150 mg Capsules are hard capsules coloured white/white, with markings of 'CLIN 150' and 'Pfizer'. They are available in blister packs (aluminium foil/PVC) of 24 and 100 capsules and bottle packs (high density polyethylene or amber glass) of 24, 100 or 500 capsules.

Not all pack sizes may be marketed.

### **Marketing Authorisation Holder**

Pfizer Limited,  
Ramsgate Road,  
Sandwich,  
Kent,  
CT13 9NJ  
UK

### **Manufacturer**

Fareva Amboise,  
Zone Industrielle,  
29 route des Industries,  
37530 Pocé-sur-Cisse,  
France

### **Company Contact address:**

If you have any comments on the way this leaflet is written, please contact Medical Information at Pfizer Limited in Walton Oaks, Tadworth, Surrey.  
Tel: 01304 616161.

**This leaflet was last revised in 02/2024.**

Ref: DA 22\_0