

FRONT



430 mm

430 mm

If you forget to take a dose, take it as soon as you remember.
If it is almost time for the next dose, skip the missed dose.
Do not take a double dose to make up for a forgotten dose.

Do not stop taking this medicine without talking to your doctor. In some cases, you may need to stop taking it gradually.

Like all medicines, this medicine can cause side effects, although not everybody gets them.

If you have an allergic reaction, see a doctor straight away. The signs may include raised lumps on your skin (weals), or swelling of your face, lips, mouth, tongue or throat.

Common (may affect up to 1 in 10 people)

- Uncommon (may affect up to 1 in 100 people)**

- Disturbed sleep.

Your doctor may take blood samples every so often to check whether Atenolol has had any effect on your blood.

- Keep this medicine out of the sight and reach of children.

Do not store above 25°C. Store your tablets in the original package.

Keep the blister strip in the carton. This will protect your medicine from light and moisture.

These measures will help protect the environment.

- The other ingredients are heavy magnesium carbonate (E504), maize starch, sodium starch glycolate (Type A), colloidal silicon dioxide, sodium lauryl sulphate, magnesium stearate (E470b), polyethylene glycol 600, hydroxypropylmethyl cellulose (E464), titanium dioxide (E171), purified talc (E553b).

They are supplied in blister pack of 14 tablets. Pack size: 14 tablets, 28 tablets, 56 tablets, 84 tablets.

Not all pack sizes may be marketed.

Fax: + 44 (0) 1915166526
Email: regulatoryfrm.mumbai@ipca.com

This leaflet was revised in 03/2022

XXXXXX

[BACK](#)

Artwork Size: 430 x 140 mm



BLACK